



































Daily Dosing of Antiretroviral Agents

	CLASS/DRUG	USUAL ADULT DAILY DOSING ^{†*}	SPECIAL CONSIDERATIONS
NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NNRTI)	VIRAMUNE [®] (nevirapine)	1 x 200 mg tablet 2 times a day 	Starting Dose: 1 x 200 mg tablet once daily for first 14 days.
	Rescriptor [®] (delavirdine mesylate)	4 x 100 mg tablets 3 times a day 	Take at least 1 hour away from antacids.
	Stocrin [®] (efavirenz)	1 x 600 mg tablet once daily 	Take at bedtime. Take away from meals.
	Intelence [®] (etravirine)	2 x 100 mg tablets 2 times a day 	Take after a meal.
NUCLEOSIDE / NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS (NRTI / NRTI)	Zerit [®] (stavudine-also known as d4T)	1 x 40 mg capsule 2 times a day  or 1 x 30 mg capsule 2 times a day 	
	Retrovir [®] (zidovudine-also known as AZT)	2 x 100 mg capsules 3 times a day  or 1 x 250 mg capsule 2 times a day 	
	3TC [®] (lamivudine)	1 x 150 mg tablet 2 times a day  or 1 x 300 mg tablet once daily 	
	Combivir [®] (lamivudine/zidovudine)	1 x 150mg / 300 mg tablet 2 times a day 	
	Ziagen [®] (abacavir)	1 x 300 mg tablet 2 times a day  or 2 x 300 mg tablets once daily 	
	Kivexa [®] (abacavir/lamivudine)	1 x 600 mg / 300 mg tablet once daily 	
	Trizivir [®] (abacavir/lamivudine/zidovudine)	1 x 300 mg / 150 mg / 300 mg tablet 2 times a day 	
	Videx [®] EC (didanosine-also known as ddl)	1 x 400 mg capsule once daily  or 1 x 250 mg capsule once daily 	Take on an empty stomach.
	Emtriva [®] (emtricitabine-also known as FTC)	1 x 200 mg capsule once daily 	
	Viread [®] (tenofovir disoproxil fumarate)	1 x 300 mg tablet once daily 	Preferably taken with a meal.
	Truvada [®] (tenofovir disoproxil fumarate/ emtricitabine)	1 x 300 mg / 200 mg once daily 	Preferably taken with a meal.

Daily Dosing of Antiretroviral Agents

	CLASS/DRUG	USUAL ADULT DAILY DOSING [†] *	SPECIAL CONSIDERATIONS
INTEGRASE INHIBITORS	Isentress[®] (raltegravir)	1 x 400 mg tablet 2 times a day 	
PROTEASE INHIBITORS (PI)	Crixivan[®] (indinavir)	2 x 400 mg capsules 3 times a day 	Take on an empty stomach. Drink at least 1.5 L of liquid daily. Lower daily doses may be used with other PIs.
	Invirase[®] (saquinavir)	2 x 500 mg tablets 2 times a day 	Take each 2 x 500 mg dose with 100 mg ritonavir. Take within 2 hours of a full meal.
	Kaletra[®] (lopinavir/ritonavir)	2 x 200 mg / 50 mg tablets 2 times a day 	Can be taken with or without food.
	Telzir[®] (fosamprenavir calcium)	1 x 700 mg tablet 2 times a day  or 2 x 700 mg tablets once daily (Not indicated for PI experienced patients) 	Take each tablet with 100 mg ritonavir. Can be taken with or without food.
	Reyataz[®] (atazanavir)	1 x 300 mg capsule once daily (Reyataz is also available in 150 mg capsules)  or 2 x 200 mg capsules once daily 	Take each 1 x 300 mg dose with 100 mg ritonavir. Take with food.
	APTIVUS[®] (tipranavir)	2 x 250 mg capsules 2 times a day 	Take each 2 x 250 mg dose with 200 mg ritonavir. Take with food.
	Prezista[®] (darunavir)	2 x 300 mg tablet 2 times a day 	Take each 2 x 300 mg dose with 100 mg ritonavir. Take with food.
	Norvir[®] (ritonavir)	Mainly used in small doses (100 mg / 200 mg) in combination with other PIs	Take with food.
FUSION AND ENTRY INHIBITORS	Fuzeon[®] (enfuvirtide-also known as T-20)	1 x 90 mg vial 2 times a day 	Give by subcutaneous injection.
	Celsentri[®] (maraviroc)	1 x 150 mg tab or 1 x 300 mg tab or 2 x 300 mg tabs 2 times a day  or  or 	Dose is dependent on concomitant medication.